HOW TO TAKE A BATHROOM BREAK ON NXSTAGE

NOTE: This information was provided by a NxStage user. The non-profit Medical Education Institute and Home Dialysis Central are not responsible for this content. Please have YOUR care team review it before use.

Steps to Stop Treatment	
	Remove tape from your shirt.
	Press STOP on the machine.
	CLAMP the arterial (RED) patient line AND the access line. DISCONNECT.
	CONNECT the arterial (RED) to the RED clamp of the saline spike.
	Keep the venous (BLUE) lines connected. Keep the clamps open.
	Do a RINSEBACK by pushing the FLUID button. CLOSE ALL CLAMPS.
	Remove the arterial (RED) line from the priming spike line.
	Remove the venous (BLUE) line from the needle.
	Connect the arterial (RED) and venous (BLUE) lines on the machine TO EACH
	OTHER using a peanut.
	Press GREEN TREATMENT Kidney on the machine. Blood will circulate in a closed
	loop through the peanut while disconnected.
	Tape your lines with syringes to your shoulder.
	hile in the bathroom, recirculate blood for a MAXIMUM of 30 minutes: Make sure RED and BLUE clamps are OPEN where connected to the peanut. Press GREEN TREATMENT Kidney on the machine and adjust the flow rates: Dialysate to 0.1 (GREEN) Ultrafiltration to 0.0 (YELLOW) Blood flow to 200 (RED)
Ste	eps to Restart Treatment
	Press STOP on machine.
	CLAMP and DISCONNECT the arterial (RED) and the venous (BLUE) patient lines
	from the peanut.
	RECONNECT the patient lines to the vascular access (RED and BLUE).
	UNCLAMP all four clamps.
	UNCLAMP RED and BLUE clamps
	Press GREEN TREATMENT Kidney on machine to resume.
	Reenter therapy rates: (RED, YELLOW, GREEN)
htt	ps://www.nxstage.com/hcp/training-resources/temporary-disconnect-with-blood-module